

This is a repository copy of *Evaluating the Impact of a Critical Time Intervention Adaptation on Health Care Utilization among Homeless Adults with Mental Health Needs in a Large Urban Center*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/171559/>

Version: Accepted Version

Article:

Reid, Nadine, Mason, Joyce, Kurdyak, Paul et al. (4 more authors) (2021) Evaluating the Impact of a Critical Time Intervention Adaptation on Health Care Utilization among Homeless Adults with Mental Health Needs in a Large Urban Center. Canadian journal of psychiatry-*Revue canadienne de psychiatrie*. ISSN 0706-7437

<https://doi.org/10.1177/0706743721996114>

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



Evaluating the impact of a Critical Time Intervention adaptation on health care utilization among homeless adults with mental health needs in a large urban centre

Journal:	<i>The Canadian Journal of Psychiatry/La Revue canadienne de psychiatrie</i>
Manuscript ID	CJP-2020-378-OR.R1
Manuscript Type:	Original Research
Date Submitted by the Author:	05-Jan-2021
Complete List of Authors:	Reid, Nadine; Centre for Addiction and Mental Health Mason, Joyce; Centre for Addiction and Mental Health; Institute for Clinical Evaluative Sciences; University of Toronto, Institute of Health Policy, Management and Evaluation Kurdyak, Paul; Centre for Addiction and Mental Health, Institute for Mental Health Policy Research ; Institute for Clinical Evaluative Sciences, Mental Health and Addictions Research Program; University of Toronto, Institute of Health Policy, Management and Evaluation; University of Toronto Faculty of Medicine, Psychiatry Nisenbaum, Rosane; St Michael's Hospital, Li Ka Shing Knowledge Institute; University of Toronto, Dalla Lana School of Public Health de Oliveira, Claire; Centre for Addiction and Mental Health, Institute for Mental Health Policy Research; Institute for Clinical Evaluative Sciences, Mental Health and Addictions Research Program; University of York, Centre for Health Economics and Hull York Medical School; University of Toronto, Institute of Health Policy, Management and Evaluation Hwang, Stephen; St Michael's Hospital, MAP Centre for Urban Health Solutions, Li Ka Shing Knowledge Institute; Institute for Clinical Evaluative Sciences, Populations & Public Health Research Program; University of Toronto Faculty of Medicine, Psychiatry; University of Toronto, Dalla Lana School of Public Health Stergiopoulos, Vicky; Centre for Addiction and Mental Health; St Michael's Hospital, MAP Centre for Urban Health Solutions, Li Ka Shing Knowledge Institute; University of Toronto Faculty of Medicine, Psychiatry
Key Words:	Critical Time Intervention, Case management, Homeless, Mental Health Services, Addictions, Healthcare utilization
Abstract:	Objective: This study evaluated the impact of a Critical Time Intervention (CTI) adaptation on health care utilization outcomes among adults experiencing homelessness and mental health needs in a large urban centre. Method: Provincial population-based administrative data from Ontario, Canada were used in a pre-post design for a cohort of 197 individuals who received the intervention between January 2013 and May 2014 and were matched to a cohort of adults experiencing homelessness who did not receive the intervention over the same time period. Changes in

	<p>health care utilization outcomes in the year pre- and post-intervention were evaluated using generalized estimating equations and post-hoc analyses evaluated differences between groups.</p> <p>Results: Pre-post analyses revealed statistically significant changes in health care utilization patterns among intervention recipients, including reduced inpatient service use and increased outpatient service use in the year following the intervention compared to the year prior. However, the matched cohort analysis found non-significant differences in health service use changes between a subgroup of intervention recipients and their matched counterparts.</p> <p>Conclusions: An adapted CTI model was associated with changes in health care utilization among people experiencing homelessness and mental health needs. However, changes were not different from those observed in a matched cohort. Rigorous study designs with adequate samples are needed to examine the effectiveness of CTI and local adaptations in diverse health care contexts.</p> <p>Abstract.docx</p>

SCHOLARONE™
Manuscripts

Title

Evaluating the impact of a Critical Time Intervention adaptation on health care utilization among homeless adults with mental health needs in a large urban centre

Authors

Nadine Reid, PhD, Project Scientist, Centre for Addiction and Mental Health

Joyce Mason, MSc, Research Methods Specialist, Centre for Addiction and Mental Health;

ICES; Institute of Health Policy, Management and Evaluation, University of Toronto

Paul Kurdyak, MD, PhD, Director, Health Outcomes and Performance Evaluation, Institute

for Mental Health Policy Research and Medical Director, Performance Improvement,

Centre for Addiction and Mental Health; Senior Core Scientist, Program Lead, Mental

Health and Addictions Research Program, ICES; Associate Professor, Department of

Psychiatry and Institute of Health Policy, Management and Evaluation, University of

Toronto

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Rosane Nisenbaum, PhD, Associate Scientist, Li Ka Shing Knowledge Institute, St. Michael’s Hospital, Unity Health Toronto; Assistant Professor, Dalla Lana School of Public Health, University of Toronto

Claire de Oliveira, PhD, Independent Scientist and Health Economist, Institute for Mental Health Policy Research, Centre for Addiction and Mental Health; Adjunct Scientist, Mental Health and Addictions Research Program, ICES; Reader in Health Economics, Centre for Health Economics and Hull York Medical School, University of York; Assistant Professor, Institute of Health Policy Management and Evaluation, University of Toronto

Stephen Hwang, MD, MPH, Director, MAP Centre for Urban Health Solutions, Li Ka Shing Knowledge Institute, St. Michael’s Hospital, Unity Health Toronto; Senior Adjunct Scientist, Populations & Public Health Research Program, ICES; Professor, Department of Psychiatry and Dalla Lana School of Public Health, University of Toronto

Vicky Stergiopoulos, MD, MHSc, Physician-in-Chief and Clinician Scientist, Centre for Addiction and Mental Health; Associate Scientist, MAP Centre for Urban Health Solutions, Li Ka Shing Knowledge Institute, St. Michael’s Hospital, Unity Health Toronto; Professor and Vice-Chair, Clinical and Innovation, Department of Psychiatry, University of Toronto

Corresponding author

Vicky Stergiopoulos

Centre for Addiction and Mental Health

100 Stokes Street, Room 6215

Toronto, Ontario M6J 1H4 Canada

Phone: 416-535-8501 Ext. 34739

Email: vicky.stergiopoulos@camh.ca

Abstract

Objective: This study evaluated the impact of a Critical Time Intervention (CTI) adaptation on health care utilization outcomes among ~~homeless~~-adults experiencing homelessness and with mental ~~illness~~-health needs in a large urban centre.

Method: Provincial population-based administrative data from Ontario, Canada were used in a pre-post design for a cohort of 197 individuals who received the intervention between January 2013 and May 2014 and were matched to a cohort of ~~homeless~~-adults experiencing homelessness who did not receive the intervention over the same time period. Changes in health care utilization outcomes in the year pre- and post-intervention were evaluated using generalized estimating equations and post-hoc analyses evaluated differences between groups.

Results: Pre-post analyses revealed statistically significant changes in health care utilization patterns among intervention recipients, including reduced inpatient service use and increased outpatient service use in the year following the intervention compared to the year prior. However, the matched cohort analysis found non-significant differences in health service use changes between a subgroup of intervention recipients and their matched counterparts.

Conclusions: An adapted CTI improved-model was associated with changes in health care utilization among homeless-people- experiencing homelessness and with mental illnesshealth needs. However, changes were not different from those observed in a matched cohort. Rigorous study designs with adequate samples are needed to examine the effectiveness of CTI and local adaptations in diverse health care contexts.

Keywords: Critical Time Intervention, case management, homeless, mental health services, addiction, health care utilization

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1. Introduction

Internationally, rates of homelessness continue to rise^{1,2}; and in Canada, national data show an increase in demand for shelter beds over time, with at least 235,000 Canadians experiencing homelessness in a year³. Homeless individualsPeople experiencing homelessness also experience higher rates of mental health and addiction challenges⁴, neurocognitive impairment^{5,6,3}, physical health problems and premature mortality^{7,84}, compared to their housed counterparts. Despite the health disparities, this population generally has limited access to appropriate, high quality health care and supports, even in settings with universal health insurance⁹⁶⁻¹²⁹. Together, these factors contribute to high rates of hospital service use and costs¹³⁴⁰⁻¹⁶⁴³.

~~With rising rates of homelessness in Canada and internationally¹⁴⁻¹⁶, Given~~
increasing rates of homelessness and the impact of homelessness on health¹⁷, t~~t~~here is an urgent need for interventions to improve health outcomes in this population, including avoidable hospital service utilization¹⁵⁴². ~~Fo~~For adults experiencing ~~homelessness and~~ mental illness, care following discharge from hospital for a mental health condition is reportedly the most important factor in reducing reliance on subsequent inpatient care¹⁸⁷. ~~Yet this population~~Adults experiencing mental illness and homelessness, however, are is

less likely to be referred to and access community-based services upon discharge from hospital, compared to housed individuals^{198,2019}.

Among ~~homeless populations~~ people experiencing homelessness, strategies to improve continuity of care by offering intensive time-limited services to support the transition from hospital to community-based services have shown promising findings²¹⁹⁻²⁶⁵. Critical time interventions (CTI) was designed to support homeless individuals during transitions of care, including the period post-hospital discharge in which people experiencing mental illness are at high risk of experiencing first-episode or recurrent homelessness in particular, offering.^{27,28} Critical time interventions involve time-limited intensive case management over a period of six to nine months and aim to help service users navigate the complex service system and establish (or re-establish) access to longer-term community-based connections, resources and interventions. This particular model, ~~has~~ ve been shown to decrease early psychiatric readmission rates^{243,296,3027}, improve perceived quality of care^{221,254}, ~~and~~ improve health and quality of life outcomes^{243,3027}, and reduce rates of homelessness^{27,31}. Past evaluations of CTIs have also suggested that this approach is cost-effective³²²⁸ in supporting transitions of care.

More evidence of the impact of CTI models, and their adaptations in diverse settings, is urgently needed to support implementation of effective and cost-effective

strategies to improve health and housing outcomes among people experiencing homelessness and who have unmet mental health needs. A recent pre-post evaluation of a CTI adaptation for ~~homeless adults~~adults experiencing homelessness and mental health needs who were transitioning from hospital to community services in Toronto, Canada, reported significant improvements in mental and physical health, substance use, and quality of life in the six months following the intervention²⁴³. Expanding on this work, the primary objective of this study was to use administrative health care utilization data to evaluate whether a brief CTI adaptation for ~~homeless adults~~adults experiencing homelessness and mental health needs who were discharged from hospital services was associated with significant ~~improvements~~changes in health care utilization outcomes in the year post-intervention compared to the year prior. This analysis in addition compared~~s~~ changes in health care utilization among participants with a recent hospitalization to those of a matched cohort of people experiencing homelessness over the same time period who did not receive the intervention.

2. Methods

2.1 Intervention

The Coordinated Access to Care for Homeless adults (CATCH-Homeless) program is a CTI adaptation in Toronto, Canada, supporting adults experiencing homelessness and

mental health needs following discharge from hospital. The program is facilitated through partnerships between three local hospitals serving large numbers of people experiencing homelessness, a primary care team, a homeless shelter, a large community mental health agency, and a physician practice plan. The program accepts referrals from all partner hospital emergency departments (EDs) and inpatient units, or from community agencies, and connects participants with transitional case managers who coordinate access to a full range of post-discharge community-based services, including mental health and addiction services, medical care, peer support, housing assistance, and other resources described in depth elsewhere³³²⁹.

2.2 Study design

This study is part of a larger mixed-methods evaluation described elsewhere^{221,3329}. Using provincial population-based administrative databases at ICES (formerly known as the Institute for Clinical Evaluative Sciences), a single-arm pre-post analysis among intervention participants was conducted to evaluate health care utilization outcomes in the year following enrolment in the intervention compared to the year prior. In addition, changes in health service utilization of a subgroup of intervention participants with a recent hospitalization were compared to those of a matched cohort of people-adults experiencing

homelessness with at least one psychiatric hospitalization over the same time period who did not receive the intervention.

2.3 Data sources

All administrative health care data were obtained from ICES. ICES is an independent, non-profit research institute funded by an annual grant from the Ontario Ministry of Health (MOH). As a prescribed entity under Ontario’s privacy legislation, ICES is authorized to collect and use health care data for the purposes of health system analysis, evaluation and decision support. Secure access to these data is governed by policies and procedures that are approved by the Information and Privacy Commissioner of Ontario. The Ontario Mental Health Reporting System (OMHRS) and the Canadian Institute for Health Information’s Discharge Abstract Database (DAD) include data on all psychiatric and acute hospitalizations, respectively. The Ontario Health Insurance Plan (OHIP) claims database contains physician billings data, and the National Ambulatory Care Reporting System gathers data on hospital- and community-based ambulatory care, such as ED visits. The Ontario Registered Persons Database (ORPD) is a registry of all individuals living in Ontario who are eligible for public health care insurance and holds data on patient demographics, such as age, sex, and postal code.

2.4 Sample

2.4.1 Intervention participants

Intervention participants were those referred to the intervention by partner sites following discharge from EDs and inpatient units. Intervention eligibility criteria included current homelessness status (defined as living in a crisis/emergency shelter, living on the street, or couch surfing), provider-determined unmet mental health needs, and participant-identified need for support services. Individuals were ineligible for Participants were excluded from the program this community-based intervention if they had recent a history of severe aggression or required severe aggression or illness severity that required residential-institutional care. Study eligibility criteria included being a resident of the province of Ontario, age 18 years and older, and not having previously received services from the program. For the current analysis, 197 individuals, who received the intervention between an index date of January 7, 2013 and May 16, 2014, completed a baseline interview, and agreed to health care record linkage were included. Twenty-six individuals (11.7% of total participants enrolled in the intervention) did not consent to health care record linkage and were excluded from the current analysis. These individuals were not different from consenting individuals in terms of gender but were younger (mean age: 34.9 years vs. 40.3 years; $P=.021$). Furthermore, they were not different in terms of baseline ED visits but had fewer baseline hospitalizations (mean: 0.7 vs. 1.2; $P=.022$). –Additional

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

~~e~~Exclusion criteria for this analysis included missing patient identifier (required for data linkage), not being enrolled in OHIP, and/or missing data on sex and/or age.

2.4.2 Matched cohort participants

The comparison group ~~obtained from the ORPD~~ was composed of individuals aged 18 years and older with recorded homelessness (as identified by a residence variable in the OMHRS or a homelessness variable in the DAD), who had at least one hospitalization for mental health or substance use during the exposure period (index date: January 1, 2013 to May 31, 2014). ~~These individuals were then matched to the intervention participants who had at least one hospitalization in the 12 months prior to their index date. Individuals who received the intervention and who had at least one hospitalization in the 12 months prior to the index date were then matched 1:2 with the comparison group of adults experiencing homelessness who had not received the intervention using propensity score-based matching. Variables used to calculate the propensity score included having a hospitalization in the 12 months prior to the index date, age, sex, neighbourhood income quintile and administrative health region of residence. We selected the closest control that met the following criteria: age within two years at the index date, same sex (hard match) and a propensity score within a caliper width of 0.20. To evaluate matching success, standardized baseline differences were calculated between the subset of intervention~~

participants and the matched cohort. Standardized differences of 0.10 or less are considered negligible³⁴. Individuals were excluded from the matched group if they were ineligible for OHIP and/or had missing sex and/or age data.

2.5 Outcome measures

To evaluate changes in acute health care utilization, count outcomes including the total number of all-cause and mental health and substance use-specific ED visits, hospitalizations, and total hospital days were calculated, ~~in addition to~~. Dichotomized outcomes indicating whether or not a participant had any all-cause and mental health and substance use-specific ED visits and hospitalizations during the observation period were also calculated. To evaluate changes in outpatient service use, count outcomes including the total number of psychiatrist visits, all-cause and mental health ~~and~~ and addiction substance use-specific general practitioner (GP) visits, and the sum total of outpatient visits were calculated, as well as dichotomized outcomes indicating whether or not a participant had any psychiatrist visits and all-cause and mental health ~~and~~ and addiction substance use-specific GP visits during the observation period. Diagnostic codes used to determine mental health and ~~addiction~~ substance use-specific hospital and outpatient visits are provided in Supplemental Table 14.

2.6 Data collection

Intervention participants' demographic and administrative health care records, held by ICES, were linked using unique encoded identifiers; deterministic linkage was applied using OHIP numbers. Health care records were examined for the 12 months pre- and post-index date. The index date for exposure was the intervention enrollment date, ranging from January 7, 2013 to May 16, 2014; the study period was January 2012 to May 2015. All participants provided written informed consent and the study received Research Ethics Board approval from Unity Health Toronto. To identify the matched cohort used in comparative analyses, population-based administrative demographic and health records were retrieved and reviewed for the same time periods.

2.7 Statistical analyses

2.7.1 Single arm pre-post analysis

Baseline characteristics for both intervention participants and individuals included in the matched cohort were calculated using frequencies and proportions for categorical variables and means and standard deviation for continuous variables. In the pre-post evaluation, inferential analyses for each count outcome included rates and rate ratios and their respective 95% confidence intervals estimated using generalized estimating equations (GEE) models with a negative binomial distribution and a log link. For each binary outcome, predicted probabilities and prevalence ratios and their respective 95% confidence

intervals were estimated using GEE models with a Poisson distribution and a log link. All models were adjusted for age, sex and neighbourhood income quintile (measured at the Census tract level). An offset variable for person-years was included to adjust for different follow-up lengths.

All analyses were conducted using SAS Enterprise Guide 7.1. A significance level of .05 was used to interpret statistical significance.

2.7.2 Matched cohort selection

———~~Individuals who received the intervention and who had at least one hospitalization in the 12 months prior to the index date were matched 1:2 with homeless individuals who had not received the intervention using propensity score-based matching. Variables used to calculate the propensity score included having a hospitalization in the 12 months prior to the index date, age, sex, neighbourhood income quintile and administrative health region of residence. We selected the closest control that met the following criteria: age within two years at the index date, same sex (hard match) and a propensity score within a caliper width of 0.20. To evaluate matching success, standardized baseline differences were calculated between the subset of intervention participants and the matched cohort. Standardized differences of 0.10 or less are considered negligible³⁰.~~

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

2.7.2 *Post-hoc comparative analysis*

Baseline demographic characteristics were calculated between intervention participants with at least one hospitalization in the 12 months pre-index date and their matched counterparts using frequencies and proportions for categorical variables and means and standard deviation for continuous variables. To show trends between groups and over time, GEE models specifying a negative binomial distribution and a log link were conducted for each count outcome to estimate rate ratios and relative rate ratios; for each binary outcome, GEE models specifying a Poisson distribution and a log link were conducted to calculate prevalence ratios and relative prevalence ratios. All post-hoc models were adjusted for age, sex and neighbourhood income quintiles and included group, time and group by time interaction variables.

3. Results

The cohort selection is detailed in Figure 1. Of the 197 eligible individuals who received the intervention during the study period, 11 health records were unlinkable (six were blank and five were invalid), resulting in a cohort of 186 participants. Of those, 51 individuals were excluded from the matched analysis because they did not have a hospitalization for mental health or substance use in the 12 months prior to the index date required for comparison with the matched sample, leaving a subset of 125 individuals.

Among 3,555 administrative records of ~~homeless individuals~~ people experiencing homelessness with a mental health or substance use-specific hospitalization during the exposure period, 250 individuals (1:2 matching) were selected for inclusion in the matched cohort.

3.1 Demographic characteristics

The demographic characteristics of intervention participants (n=186) are presented in Supplemental Table 24. The majority of the sample (78.5%) was male. Participants had a mean age of 40.3 years ($SD=12.0$), with an even distribution across age brackets between 18 and 59 years; only 4.3% of participants were over the age of 60.

Table 1 shows the demographic characteristics of the 125 intervention participants included in the matched cohort analysis and their matched counterparts; these were similar to those of the full cohort of intervention participants in age, sex, and neighbourhood income quintile as indicated by no standardized difference between the two groups being greater than 0.10.

3.2 Pre-post analysis

Rates per person-year and rate ratios (RR) for intervention participants (n=186) in the 12 months pre- and post-intervention enrolment for the number of hospital admissions,

hospital days, ED visits and outpatient visits are presented in Table 2; results for binary outcomes are presented in Supplemental Table 32. Overall, results reveal changes in health care utilization patterns, with decreased inpatient care and increased outpatient care in the 12 months following enrolment in the intervention, compared to the 12 months prior. Inpatient hospital use decreased significantly among intervention participants in the 12 months following enrolment, especially for mental health and substance use-specific visits. Adjusted admission rates decreased significantly for both all cause and mental health and substance use--specific causes, by 33% (95% CI 15%-47%; $P=.001$) and 43% (95% CI 26%-66%; $P<.001$), respectively, in the 12 months post-intervention relative to 12 months prior. The total number of hospital days also decreased significantly, by 54% (95% CI 36%-77%; $P<.001$) for all-cause days and by 63% (95% CI 48%-74%; $P<.001$) for mental health and substance use days. While the number of ED visits did not change significantly over time, the overall prevalence of ED visits did decrease significantly, by 14% for any reason (95% CI 7%-20%; $P<.001$); and by 31% for mental health and substance use reasons (95% CI 12%-39%; $P<.001$) (Supplemental Table 32). In contrast to the decrease in acute hospital service use, outpatient psychiatrist visits increased by 40% (95% CI 1.15-1.70; $P=.001$) in the 12 months following the intervention compared to the 12 months prior; GP visits did not change significantly.

3.3 Matched cohort analysis

Rate ratios and relative rate ratios (RRR) for the subset of intervention participants with at least one hospitalization in the 12 months prior to the index date (n=125) compared to matched controls (n=250) 12 months pre- and post-intervention, for the number of hospital admissions, hospital days, ED visits and outpatient visits are presented in Table 3. ³⁵ Prevalence ratios (PR) and relative prevalence ratios (RPR) are presented in Supplemental Table 4. ³⁶ For both count and binary outcomes, results of the matched cohort analysis suggest that while results generally trended in the same direction as the pre-post analysis, the changes in health service use experienced by intervention participants in the year following receipt of the intervention as compared to the year prior were not significantly different than the changes experienced by their matched counterparts (Table 3).

4. Discussion

Care continuity ^{35,36} is essential to the delivery of high quality services for people living with chronic health conditions, including mental illness and addictions, and is associated with improved health and service use outcomes ^{36,2}. Our findings of decreased acute care service use and increased outpatient service use in the 12 months following the intervention as compared to the 12 months prior are consistent with previous outcomes of critical time intervention (CTI) studies, finding significant ~~improvements~~ changes in

outcome measures^{243,296,3027}. In addition, our findings further support and are supported by previous qualitative research suggesting improved experiences of continuity of care among ~~homeless adults~~adults experiencing homelessness receiving CTI or CTI adaptations^{224,373}. Notably, our findings highlight continued high rates of emergency department (ED) visits post-intervention among participants, suggesting that this population may have more acute and/or enduring needs than the brief CTI adaptation can address in our local context. Access to housing and high quality intensive case management in Toronto, for example, are extremely limited and hindered by long wait lists; it is possible that participants continued to visit the ED for immediate shelter and support post-intervention, as intervention resources were limited and not linked to housing. Persisting high ED use among participants reaffirms the need for inclusion of housing in mental health policy priorities.¹⁷

Although prior controlled studies of CTI have demonstrated improvements in acute care utilization in some settings²⁶⁵, the addition of a matched cohort analysis in the current study indicates that changes in service use patterns of recipients of a brief CTI adaptation within our setting of universal health insurance and a relatively service-rich environment are not significantly different than those of their matched counterparts.

Similar to this analysis, a prior post-hoc analysis of intervention participants using self-reported data and a comparison group of ~~homeless adults~~adults experiencing

~~homelessness with and mental illness-mental illness who were~~ receiving usual care found successful reduction in mental health symptoms and alcohol and drug use problems among intervention participants, but increased ED visits and days spent in hospital over six months²⁴³. The current ~~post-hoc~~ analysis, ~~in~~ using administrative data and a cohort matched on prior hospitalizations, ~~adds-was intended to build off the limitations of the~~ ~~aforementioned prior analysis~~²⁴ ~~by focusing on service use outcomes using administrative data and exploring between-group differences with added methodological~~ ~~additional~~ rigor and identifies no significant difference in service use patterns among intervention and matched cohort participants. ~~Additional measures including clinical characteristics such as diagnosis and acuity, and demographic and service use factors such as geographical location and resource availability, may help to better explain the identified non-significant differences between groups in future studies.~~

Our findings further ~~highlight~~ the need for rigorous methods in evaluating new interventions in diverse contexts. ~~Furthermore, it points to the need to ensure fidelity to key ingredients when adapting evidence-supported interventions in diverse contexts.~~ The ~~brief~~ CTI adaptation evaluated in the current study was set in a large urban centre under a universal health insurance system and included a dedicated, low-barrier, multidisciplinary weekly ~~physician~~ clinic in addition to brief case management support with limited training

and supervision of frontline staff, in keeping with a pragmatic field intervention. Our findings suggest that participants’ access to preventative outpatient services ~~s-use~~ was successfully facilitated by ~~appropriately designed and dedicated~~ streamlined access to GPs and psychiatrists. Such an approach may be particularly relevant to similarly large urban centre where timely access to ~~appropriate physician and case management~~ resources is hindered by rapid general and homeless population growth and corresponding demand for services.

While fidelity standards for CTI have been developed, detailing the key model components requiring adherence, and the contextual structures and staff competencies needed to ensure model integrity³⁸⁴⁻⁴⁰³⁶, a recent systematic review indicates that fidelity and adaptations remain highly variable²⁶⁵ and should be a required component of rigorous evaluations. The fidelity of the intervention to the CTI model was not formally assessed in this study. Future efforts should ensure that local adaptations of evidence-based interventions balance local needs and resources with fidelity to key intervention ingredients, so that evidence generated can be reliably attributed to the model of interest.

4.1 Strengths and limitations

Our findings contribute to the growing evidence on the impact of CTI and its adaptations on health service use. Results are strengthened by the study’s methodological

approach, which included the use of population-based administrative data and a rigorous selection process to ensure the appropriateness of comparators. However, the identification of a matched cohort was challenged by the lack of a baseline acuity measure, requiring us to match on recent hospitalizations as a proxy for acuity, although post-matching demographic comparisons suggested the groups were minimally different. Additionally, intervention participants were selected using some clinical criteria that could not be applied to matched individuals identified from administrative data. In the absence of a built-in control group, this an acknowledged design limitation that could have resulted in selection bias.

~~Additionally, we~~We were also limited by our sample size and underpowered to detect hospitalization differences. It is possible that the lower relative rates of psychiatric hospitalizations observed in the intervention group would be significantly different with a larger sample. Still, without a randomized design, it is possible the improvements in health care use observed in the pre-post analysis are due to regression to the mean, rather than the effect of the intervention.

Given the high cost of hospitalizations and from a quality of care perspective, future research should focus on rigorous evaluations using experimental methods with large samples and robust sets of explanatory variables, in addition to approaches to strengthening

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

fidelity of local adaptations to evidence-supported treatments. Related research should investigate the cost-effectiveness of these interventions to maximize health outcomes given the limited resources available.

5. Conclusions

Study findings suggest that participants of a brief Critical Time Intervention (CTI) for adults experiencing homelessness and unmet mental health needs had decreased acute care use and increased outpatient service use post-intervention. Post-hoc analyses, however, found that changes in service use patterns were not significantly different from those of a matched cohort of adults experiencing homelessness. While ~~Critical Time Intervention-CTI~~ and its adaptations hold promise in improving continuity of care and health outcomes among homeless people with mental illness for this population, more r. Rigorous study designs with adequate sample sizes are needed to further examine the effectiveness of CTI and local adaptations in diverse health care contexts.

Data Access

Data included in this publication are not publicly available and were collected by ICES. As a prescribed entity under Ontario's privacy legislation, ICES is authorized to collect and use health care data for the purposes of health system analysis, evaluation and decision support. Secure access to these data is governed by policies and procedures that are approved by the Information and Privacy Commissioner of Ontario. For further questions regarding data accessibility, please contact the corresponding author.

Acknowledgements

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

This study was supported by ICES, which is funded by an annual grant from the Ontario Ministry of Health (MOH). Parts of this material are based on data and information compiled and provided by: MOH and the Canadian Institute for Health Information. The analyses, conclusions, opinions and statements expressed herein are solely those of the authors and do not reflect those of the funding or data sources; no endorsement is intended or should be inferred.

Conflict of Interest Disclosure: none.

Funding: This study was supported by ICES, which is funded by an annual grant from the Ontario Ministry of Health.

References

1. [Abbé Pierre Foundation, European Federation of National Organisations Working with the Homeless \[FEANTSA\]. Fifth Overview of Housing Exclusion in Europe 2020. FEANTSA: Brussels; 2020.](#)
2. [Henry M, Watt R, Mahathey A et al. The 2019 Annual Homeless Assessment Report \(AHAR\) to Congress: Part 1. Point-in-time estimates of homelessness. The U.S. Department of Housing and Urban Development; 2020.](#)

3. Gaetz S, DeJ E, Richter T et al. The State of Homelessness in Canada 2016.

Toronto: Canadian Observatory on Homelessness Press; 2016.

4.4. Fazel S, Geddes JR, Kushel M. The health of homeless people in high-income countries: Descriptive epidemiology, health consequences, and clinical and policy recommendations. *Lancet*. 2014;384(9953):1529-1540.

[https://doi.org/10.1016/S0140-6736\(14\)61132-6](https://doi.org/10.1016/S0140-6736(14)61132-6)

4.5. Depp CA, Vella L, Orff HJ et al. A quantitative review of cognitive functioning in homeless adults. *J Nerv Ment Dis*. 2015;203(2):126-131.

<https://doi.org/10.1097/NMD.0000000000000248>

4.6. Stergiopoulos V, Cusi A, Bekele T et al. Neurocognitive impairment in a large sample of homeless adults with mental illness. *Acta Psychiatr Scand*.

2015;131(4):256-268. <https://doi.org/10.1111/acps.12391>

4.7. Hwang SW. Homelessness and health. *CMAJ*. 2001;164(2):229-233.

5.8. Aldridge RW, Story A, Hwang SW et al. Morbidity and mortality in homeless individuals, prisoners, sex workers, and individuals with substance use disorders in high-income countries: A systematic review and meta-analysis. *Lancet*. 2018;391(10117):241-250. [https://doi.org/10.1016/S0140-6736\(17\)31869-X](https://doi.org/10.1016/S0140-6736(17)31869-X)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

~~6-9.~~ Corrigan P, Pickett S, Kraus D et al. Community-based participatory research examining the health care needs of African Americans who are homeless with mental illness. *J Health Care Poor Underserved*. 2015;26(1):119-133. <https://doi.org/10.1353/hpu.2015.0018>

~~7-10.~~ Gelberg L, Browner CH, Lejano E et al. *Women & Health*. 2004;40(2):87-100. https://doi.org/10.1300/J013v40n02_06

~~8-11.~~ Khandor E, Mason K. *The Street Health Report 2007*. Toronto, ON: Street Health; 2007.

~~9-12.~~ Skosireva A, O’Campo P, Zerger S et al. Different faces of discrimination: Perceived discrimination among homeless adults with mental illness in healthcare settings. *BMC Health Serv Res*. 2014;14:376. <https://doi.org/10.1186/1472-6963-14-376>

~~10-13.~~ Buck DS, Brown CA, Mortensen K et al. Comparing homeless and domiciled patients’ utilization of the Harris County, Texas public hospital system. *J Health Care Poor Underserved*. 2012;23(4):1660-1670. <https://doi.org/10.1353/hpu.2012.0171>

~~11-14.~~ Fazel S, Geddes JR, Kushel M. The health of homeless people in high-income countries: Descriptive epidemiology, health consequences, and clinical and

policy recommendations. *Lancet*. 2014;384(9953):1529-1540.

[https://doi.org/10.1016/S0140-6736\(14\)61132-6](https://doi.org/10.1016/S0140-6736(14)61132-6)

- ~~12-15.~~ Hwang SW, Weaver J, Aubry T et al. Hospital costs and length of stay among homeless patients admitted to medical, surgical, and psychiatric services. *Med Care*. 2011;49(4):350-354. <https://doi.org/10.1097/MLR.0b013e318206c50d>
- Saab D, Bisenbaum R, Dhalla I et al. Hospital readmissions in a community-based sample of homeless adults: A matched-cohort study. *J Gen Intern Med*. 2016;31(9):1011-8. <https://doi.org/10.1007/s11606-016-3680-8>

~~16.~~

- ~~17.~~ Pevalin DJ, Reeves A, Baker E et al. The impact of persistent poor housing conditions on mental health: A longitudinal population-based study. *Prev Med*. 2017;105:304-310. <https://doi.org/10.1016/j.ypmed.2017.09.020>

- ~~13.~~ Baggett TP, O'Connell JJ, Singer DE et al. The unmet health care needs of homeless adults: A national study. *Am J Public Health*. 2010;100(7):1326-1333. <https://doi.org/10.2105/AJPH.2009.180109>

- ~~14-18.~~ Klinkenberg WD, Calsyn RJ. Predictors of psychiatric hospitalization: A multivariate analysis. *Adm Policy Ment Health*. 1998;25(4):403-410. <https://doi.org/10.1023/A:1022292407868>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

[15-19.](#) Burra TA, Hwang SW, Rourke SB et al. Homeless and housed inpatients with schizophrenia: Disparities in service access upon discharge from hospital. *Int J Ment Health Addict.* 2012;10(5):778-789. <https://doi.org/10.1007/s11469-012-9392-3>

[16-20.](#) Laliberte V, Stergiopoulos V, Jacob B et al. Homelessness at discharge and its impact on psychiatric readmission and physician follow-up: A population-based cohort study. *Epidemiol Psychiatr Sci.* 2020;29:e21. <https://doi.org/10.1017/S2045796019000052>

[17-21.](#) Hwang SW, Burns T. Health interventions for people who are homeless. *Lancet.* 2014;384(9953):1541-1547. [https://doi.org/10.1016/S0140-6736\(14\)61133-8](https://doi.org/10.1016/S0140-6736(14)61133-8)

[18-22.](#) Lamanna D, Stergiopoulos V, Durbin J et al. Promoting continuity of care for homeless adults with unmet health needs: The role of brief interventions. *Health Soc Care Community.* 2018;26(1):56-64. <https://doi.org/10.1111/hsc.12461>

[19-23.](#) Stergiopoulos V, Gozdzik A, Misir V et al. Effectiveness of Housing First with intensive case management in an ethnically diverse sample of homeless adults with mental illness: A randomized controlled trial. *PLoS One.* 2015;10(7):e0130281. <https://doi.org/10.1371/journal.pone.0130281>

20-24. Stergiopoulos V, Gozdzik A, Nisenbaum R et al. Bridging hospital and community care for homeless adults with mental health needs: Outcomes of a brief interdisciplinary intervention. *Can J Psychiatry*. 2018;63(11):774-784.

<https://doi.org/10.1177/0706743718772539>

21-25. Tomita A, Herman DB. The role of a critical time intervention on the experience of continuity of care among persons with severe mental illness following hospital discharge. *J Nerv Ment Dis*. 2015;203(1):65-70.

<https://doi.org/10.1097/NMD.0000000000000224>

22-26. Ponka D, Agbata E, Kendall C et al. The effectiveness of case management interventions for the homeless, vulnerably housed and persons with lived experience: A systematic review. *PLoS One*. 2020;15(4):e0230896.

<https://doi.org/10.1371/journal.pone.0230896>

27. Herman DB, Conover S, Gorroochurn P et al. Randomized trial of critical time intervention to prevent homelessness after hospital discharge. *Psychiatr Serv*. 2011;62(7):713-719. https://doi.org/10.1176/ps.62.7.pss6207_0713

28. Herman DB, Susser ES, Jandorf L et al. Homelessness among individuals with psychotic disorders hospitalized for the first time: Findings from the Suffolk County

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Mental Health Project. Am J Psychiatry. 1998;155(1):109-113.
<https://doi.org/10.1176/ajp.155.1.109>

23-29. Shaffer SL, Hutchison SL, Ayers AM et al. Brief critical time intervention to reduce psychiatric rehospitalization. Psychiatr Serv. 2015;66(11):1155-1161.
<https://doi.org/10.1176/appi.ps.201400362>

30. Tomita A, Herman DB. The impact of critical time intervention in reducing psychiatric rehospitalization after hospital discharge. Psychiatr Serv. 2012;63(9):935-937. <https://doi.org/10.1176/appi.ps.201100468>

24-31. KasproW WJ, Rosenheck RA. Outcomes of critical time intervention case management of homeless veterans after psychiatric hospitalization. Psychiatr Serv. 2007;58(7):929-935. <https://doi.org/10.1176/ps.2007.58.7.929>.

25-32. Jones K, Colson PW, Holter MC. Cost-effectiveness of critical time intervention to reduce homelessness among persons with mental illness. Psychiatr Serv. 2003;54(6):884-90. <https://doi.org/10.1176/appi.ps.54.6.884>

26-33. Stergiopoulos V, Gozdzik A, Nisenbaum R et al. Integrating hospital and community care for homeless people with unmet mental health needs: Program rationale, study protocol and sample description of a brief multidisciplinary case

management intervention. *Int J Ment Health Addict*. 2017;15:362-378.

<https://doi.org/10.1007/s11469-017-9731-5>

[27-34.](#) Austin PC. Using the standardized difference to compare the prevalence of a binary variable between two groups in observational research. *Commun Stat-Simul Comput*. 2009;38(6):1228-1234. <https://doi.org/10.1080/03610910902859574>

[28-35.](#) Haggerty JL, Reid RJ, Freeman GK et al. Continuity of care: A multidisciplinary review. *Br Med J*. 2003;327(7425):1219-1221. <https://doi.org/10.1136/bmj.327.7425.1219>

[29-36.](#) Puntis S, Rugkasa J, Forrest A et al. Associations between continuity of care and patient outcomes in mental health care: A systematic review. *Psychiatr Serv*. 2015;66(4):354-363. <https://doi.org/10.1176/appi.ps.201400178>

[30-37.](#) Dixon L, Goldberg R, Iannone V et al. Use of a critical time intervention to promote continuity of care after psychiatric inpatient hospitalization. *Psychiatr Serv*. 2009;60(4):451-458. <https://doi.org/10.1176/ps.2009.60.4.451>

[31-38.](#) da Silva TFC, Lovisi GM, Conover S. Developing an instrument for assessing fidelity to the intervention in the Critical Time Intervention – Task Shifting (CTI-TS) – preliminary report. *Arch Psychiatry Psychother*. 2014;1:55-62. <https://doi.org/10.12740/APP/23279>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

[32-39.](#) De Vet R, Lako DAM, Jeijersbergen MD et al. Critical time intervention for people leaving shelters in the Netherlands: Assessing fidelity and exploring facilitators and barriers. *Adm Policy Ment Health*. 2017;44:67-80. <https://doi.org/10.1007/s10488-015-0699-9>

[33-40.](#) Shaw J, Conover S, Herman D et al. Critical time intervention for severely mentally ill prisoners (CrISP): A randomized controlled trial. *Health Services and Delivery Research*. 2017;5(8). <https://doi.org/10.3310/hsdr05080>

Tables

Table 1: Demographic characteristics of CATCH-H participants *with at least one hospitalization in the 12 months-pre-intervention* (n=125) and matched controls (n= 250)

Demographic variables		CATCH-H participants with hospitalization (n=125)		Matched controls (n=250)		Standardized difference
		n	%	n	%	
Age (years)	18 to 29	29	23.2	58	23.2	0
	30 to 39	27	21.6	52	20.8	0.02

	40 to 49	36	28.8	73	29.2	0.01
	50 to 59	25	20	49	19.6	0.01
	60 and over	8	6.4	18	7.2	0.03
	Mean (SD)	41.1 (12.4)		41	12.3	0
Sex	Female	27	21.6	52	20.8	0
	Male	98	78.4	19	79.2	
				8		
Neighbourhood income quintile	1* (low)	39	31.2	90	36	0.09
	2 (medium low)	30	24	55	22	0.05
	3 (medium)	26	20.8	49	19.6	0.03
	4 (medium high)	13	10.4	23	9.2	0.04
	5 (high)	17	13.6	33	13.2	0.01

Legend: SD = standard deviation

*1: includes individuals in the 1st income quintile and those where this information was missing

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2: Rates per person-year and rate ratios (RR) for CATCH-H participants (n=186) 12 months pre-intervention vs. 12 months post-intervention for number of hospital admissions, hospital days, ED visits and outpatient visits estimated from adjusted generalized estimating equations with negative binomial distribution

Outcome variable		CATCH-H participants (n=186)							
		Rate per person-year				Rate ratio (RR)			
		12 months pre-intervention		12 months post-intervention		RR	95% CI	P-value	
		Rate	95% CI	Rate	95% CI				
Hospital admissions	All cause	2.00	1.54-2.60	1.35	0.96-1.88	0.67	0.53-0.85	.001	
	Mental health and substance use	1.22	0.92-1.63	0.69	0.48-1.01	0.57	0.44-0.74	<.001	
Days in hospital	All cause	24.66	18.93-32.12	11.37	8.09-15.99	0.46	0.33-0.64	<.001	
	Mental health and substance use	17.67	12.34-25.32	6.47	4.22-9.93	0.37	0.26-0.52	<.001	
ED visits	All cause	8.98	6.76-11.94	9.57	6.90-13.29	1.07	0.90-1.25	.446	
	Mental health and substance use	3.77	2.79-5.08	3.57	2.49-5.11	0.95	0.75-1.19	.645	
Outpatient visits	Psychiatrist	3.16	2.36-4.25	4.42	3.32-5.88	1.40	1.15-1.70	.001	
	GP	All cause	7.49	6.30-8.91	7.70	6.63-8.94	1.03	0.88-1.20	.725
		Mental health and substance use	3.38	2.70-4.23	3.80	3.16-4.58	1.13	0.93-1.37	.224
		Total outpatient visits	11.05	9.35-13.07	12.64	10.81-14.77	1.14	1.00-1.31	.049

(Psychiatrist +
GP)

Legend: CI = confidence interval; GP = general practitioner

Table 3: Rate ratios (RR) and relative rate ratios (RRR) for *CATCH-H* participants with at least one hospitalization in the 12 months pre-intervention (n=125) compared to matched controls (n=250) 12 months pre-intervention vs. 12 months post-intervention for number of hospital admissions, hospital days, ED visits and outpatient visits estimated using adjusted generalized estimating equations with negative binomial distribution


Outcome variable		12 months pre-intervention			12 months post-intervention			Relative rate ratio (RRR) 12 months post-intervention vs. 12 months pre-intervention		
		RR	95% CI	P-value	RR	95% CI	P-value	RR	95% CI	P-value
Hospital admissions	All cause		0.99							
		1.20	1.45	.057	1.17	0.82-1.66	.38	0.97	0.71-1.33	.870
	Mental health and substance use	1.06	0.87-1.29	.565	0.95	0.64-1.39	.78	0.89	0.63-1.26	.522
Days in hospital	All cause	1.04	0.75-1.43	.808	0.77	0.51-1.17	.22	0.74	0.46-1.20	.225
	Mental health and substance use	0.93	0.67-1.29	.667	0.60	0.38-0.95	.03	0.64	0.38-1.10	.104
ED visits	All cause	1.35	1.01-1.79	.040	1.69	1.19-2.42	.004	1.26	0.95-1.67	.113

Outpatient visits	Mental health and substance use		0.95							
			-			1.11-2.43		1.23	0.89-1.71	.212
		1.33	1.86	.096	1.64		.01			
	Psychiatrist		0.96							
			-			1.02-1.77		1.04	0.77-1.40	.806
		1.29	1.74	.091	1.34					
	G P	All cause								
			-	.10		1.06-1.75		1.11	0.86-1.44	.409
		1.22	1.56	2	1.36		.02			
	Mental health and substance use		0.85							
			-	.35		0.98-1.77		1.14	0.84-1.55	.408
		1.16	1.58	6	1.32		.07			
Total outpatient visits			1.02							
(Psychiatrist + GP)			-			1.10-1.62		1.07	0.87-1.33	.522
		1.24	1.52	.035	1.33		.003			

Legend: CI = confidence interval; GP = general practitioner

Figures

Figure 1. Exposed and matched cohort selection



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Title

Evaluating the impact of a Critical Time Intervention adaptation on health care utilization among homeless adults with mental health needs in a large urban centre

Authors

Nadine Reid, PhD, Project Scientist, Centre for Addiction and Mental Health

Joyce Mason, MSc, Research Methods Specialist, Centre for Addiction and Mental Health; ICES; Institute of Health Policy, Management and Evaluation, University of Toronto

Paul Kurdyak, MD, PhD, Director, Health Outcomes and Performance Evaluation, Institute for Mental Health Policy Research and Medical Director, Performance Improvement, Centre for Addiction and Mental Health; Senior Core Scientist, Program Lead, Mental Health and Addictions Research Program, ICES; Associate Professor, Department of Psychiatry and Institute of Health Policy, Management and Evaluation, University of Toronto

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Rosane Nisenbaum, PhD, Associate Scientist, Li Ka Shing Knowledge Institute, St. Michael’s Hospital, Unity Health Toronto; Assistant Professor, Dalla Lana School of Public Health, University of Toronto

Claire de Oliveira, PhD, Independent Scientist and Health Economist, Institute for Mental Health Policy Research, Centre for Addiction and Mental Health; Adjunct Scientist, Mental Health and Addictions Research Program, ICES; Reader in Health Economics, Centre for Health Economics and Hull York Medical School, University of York; Assistant Professor, Institute of Health Policy Management and Evaluation, University of Toronto

Stephen Hwang, MD, MPH, Director, MAP Centre for Urban Health Solutions, Li Ka Shing Knowledge Institute, St. Michael’s Hospital, Unity Health Toronto; Senior Adjunct Scientist, Populations & Public Health Research Program, ICES; Professor, Department of Psychiatry and Dalla Lana School of Public Health, University of Toronto

Vicky Stergiopoulos, MD, MHSc, Physician-in-Chief and Clinician Scientist, Centre for Addiction and Mental Health; Associate Scientist, MAP Centre for Urban Health Solutions, Li Ka Shing Knowledge Institute, St. Michael’s Hospital, Unity Health Toronto; Professor and Vice-Chair, Clinical and Innovation, Department of Psychiatry, University of Toronto

Corresponding author

Vicky Stergiopoulos

Centre for Addiction and Mental Health

100 Stokes Street, Room 6215

Toronto, Ontario M6J 1H4 Canada

Phone: 416-535-8501 Ext. 34739

Email: vicky.stergiopoulos@camh.ca

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

Objective: This study evaluated the impact of a Critical Time Intervention (CTI) adaptation on health care utilization outcomes among adults experiencing homelessness and mental health needs in a large urban centre.

Method: Provincial population-based administrative data from Ontario, Canada were used in a pre-post design for a cohort of 197 individuals who received the intervention between January 2013 and May 2014 and were matched to a cohort of adults experiencing homelessness who did not receive the intervention over the same time period. Changes in health care utilization outcomes in the year pre- and post-intervention were evaluated using generalized estimating equations and post-hoc analyses evaluated differences between groups.

Results: Pre-post analyses revealed statistically significant changes in health care utilization patterns among intervention recipients, including reduced inpatient service use and increased outpatient service use in the year following the intervention compared to the year prior. However, the matched cohort analysis found non-significant differences in health service use changes between a subgroup of intervention recipients and their matched counterparts.

Conclusions: An adapted CTI model was associated with changes in health care utilization among people experiencing homelessness and mental health needs. However, changes were not different from those observed in a matched cohort. Rigorous study designs with adequate samples are needed to examine the effectiveness of CTI and local adaptations in diverse health care contexts.

Keywords: Critical Time Intervention, case management, homeless, mental health services, addiction, health care utilization

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1. Introduction

Internationally, rates of homelessness continue to rise^{1,2}; and in Canada, national data show an increase in demand for shelter beds over time, with at least 235,000 Canadians experiencing homelessness in a year³. People experiencing homelessness also experience higher rates of mental health and addiction challenges⁴, neurocognitive impairment^{5,6}, physical health problems and premature mortality^{7,8}, compared to their housed counterparts. Despite the health disparities, this population generally has limited access to appropriate, high quality health care and supports, even in settings with universal health insurance⁹⁻¹². Together, these factors contribute to high rates of hospital service use and costs¹³⁻¹⁶.

Given increasing rates of homelessness and the impact of homelessness on health¹⁷, there is an urgent need for interventions to improve health outcomes in this population, including avoidable hospital service utilization¹⁵. For adults experiencing mental illness, care following discharge from hospital for a mental health condition is reportedly the most important factor in reducing reliance on subsequent inpatient care¹⁸. Adults experiencing mental illness and homelessness, however, are less likely to be referred to and access community-based services upon discharge from hospital, compared to housed individuals^{19,20}.

Among people experiencing homelessness, strategies to improve continuity of care by offering intensive time-limited services to support the transition from hospital to community-based services have shown promising findings²¹⁻²⁶. Critical time intervention (CTI) was designed to support homeless individuals during transitions of care, including the period post-hospital discharge in which people experiencing mental illness are at high risk of experiencing first-episode or recurrent homelessness.^{27,28} Critical time interventions involve time-limited intensive case management over a period of six to nine months and aim to help service users navigate the complex service system and establish (or re-establish) access to longer-term community-based connections, resources and interventions. This particular model has been shown to decrease early psychiatric readmission rates^{24,29,30}, improve perceived quality of care^{22,25}, improve health and quality of life outcomes^{24,30}, and reduce rates of homelessness^{27,31}. Past evaluations of CTIs have also suggested that this approach is cost-effective³² in supporting transitions of care.

More evidence of the impact of CTI models, and their adaptations in diverse settings, is urgently needed to support implementation of effective and cost-effective strategies to improve health and housing outcomes among people experiencing homelessness and who have unmet mental health needs. A recent pre-post evaluation of a CTI adaptation for adults experiencing homelessness and mental health needs who were

transitioning from hospital to community services in Toronto, Canada, reported significant improvements in mental and physical health, substance use, and quality of life in the six months following the intervention²⁴. Expanding on this work, the primary objective of this study was to use administrative health care utilization data to evaluate whether a brief CTI adaptation for adults experiencing homelessness and mental health needs who were discharged from hospital services was associated with significant changes in health care utilization outcomes in the year post-intervention compared to the year prior. This analysis in addition compared changes in health care utilization among participants with a recent hospitalization to those of a matched cohort of people experiencing homelessness over the same time period who did not receive the intervention.

2. Methods

2.1 Intervention

The Coordinated Access to Care for Homeless adults (CATCH-Homeless) program is a CTI adaptation in Toronto, Canada, supporting adults experiencing homelessness and mental health needs following discharge from hospital. The program is facilitated through partnerships between three local hospitals serving large numbers of people experiencing homelessness, a primary care team, a homeless shelter, a large community mental health agency, and a physician practice plan. The program accepts referrals from all partner

hospital emergency departments (EDs) and inpatient units, or from community agencies, and connects participants with transitional case managers who coordinate access to a full range of post-discharge community-based services, including mental health and addiction services, medical care, peer support, housing assistance, and other resources described in depth elsewhere³³.

2.2 Study design

This study is part of a larger mixed-methods evaluation described elsewhere^{22,33}. Using provincial population-based administrative databases at ICES (formerly known as the Institute for Clinical Evaluative Sciences), a single-arm pre-post analysis among intervention participants was conducted to evaluate health care utilization outcomes in the year following enrolment in the intervention compared to the year prior. In addition, changes in health service utilization of a subgroup of intervention participants with a recent hospitalization were compared to those of a matched cohort of adults experiencing homelessness with at least one psychiatric hospitalization over the same time period who did not receive the intervention.

2.3 Data sources

All administrative health care data were obtained from ICES. ICES is an independent, non-profit research institute funded by an annual grant from the Ontario Ministry of Health (MOH). As a prescribed entity under Ontario’s privacy legislation, ICES is authorized to collect and use health care data for the purposes of health system analysis, evaluation and decision support. Secure access to these data is governed by policies and procedures that are approved by the Information and Privacy Commissioner of Ontario. The Ontario Mental Health Reporting System (OMHRS) and the Canadian Institute for Health Information’s Discharge Abstract Database (DAD) include data on all psychiatric and acute hospitalizations, respectively. The Ontario Health Insurance Plan (OHIP) claims database contains physician billings data, and the National Ambulatory Care Reporting System gathers data on hospital- and community-based ambulatory care, such as ED visits. The Ontario Registered Persons Database (RPDB) is a registry of all individuals living in Ontario who are eligible for public health care insurance and holds data on patient demographics, such as age, sex, and postal code.

2.4 Sample

2.4.1 Intervention participants

Intervention participants were those referred to the intervention by partner sites following discharge from EDs and inpatient units. Intervention eligibility criteria included

current homelessness status (defined as living in a crisis/emergency shelter, living on the street, or couch surfing), provider-determined unmet mental health needs, and participant-identified need for support services. Individuals were ineligible for this community-based intervention if they had recent severe aggression or illness severity that required institutional care. Study eligibility criteria included being a resident of the province of Ontario, age 18 years and older, and not having previously received services from the program. For the current analysis, 197 individuals, who received the intervention between an index date of January 7, 2013 and May 16, 2014, completed a baseline interview, and agreed to health care record linkage were included. Twenty-six individuals (11.7% of total participants enrolled in the intervention) did not consent to health care record linkage and were excluded from the current analysis. These individuals were not different from consenting individuals in terms of gender but were younger (mean age: 34.9 years vs. 40.3 years; $P=.021$). Furthermore, they were not different in terms of baseline ED visits but had fewer baseline hospitalizations (mean: 0.7 vs. 1.2; $P=.022$). Additional exclusion criteria for this analysis included missing patient identifier (required for data linkage), not being enrolled in OHIP, and/or missing data on sex and/or age.

2.4.2 Matched cohort participants

The comparison group was composed of individuals aged 18 years and older with recorded homelessness (as identified by a residence variable in the OMHRS or a homelessness variable in the DAD), who had at least one hospitalization for mental health or substance use during the exposure period (index date: January 1, 2013 to May 31, 2014). Individuals who received the intervention and who had at least one hospitalization in the 12 months prior to the index date were then matched 1:2 with the comparison group of adults experiencing homelessness who had not received the intervention using propensity score-based matching. Variables used to calculate the propensity score included having a hospitalization in the 12 months prior to the index date, age, sex, neighbourhood income quintile and administrative health region of residence. We selected the closest control that met the following criteria: age within two years at the index date, same sex (hard match) and a propensity score within a caliper width of 0.20. To evaluate matching success, standardized baseline differences were calculated between the subset of intervention participants and the matched cohort. Standardized differences of 0.10 or less are considered negligible³⁴. Individuals were excluded from the matched group if they were ineligible for OHIP and/or had missing sex and/or age data.

2.5 Outcome measures

To evaluate changes in acute health care utilization, count outcomes including the total number of all-cause and mental health and substance use-specific ED visits, hospitalizations, and total hospital days were calculated. Dichotomized outcomes indicating whether or not a participant had any all-cause and mental health and substance use-specific ED visits and hospitalizations during the observation period were also calculated. To evaluate changes in outpatient service use, count outcomes including the total number of psychiatrist visits, all-cause and mental health and substance use-specific general practitioner (GP) visits, and the sum total of outpatient visits were calculated, as well as dichotomized outcomes indicating whether or not a participant had any psychiatrist visits and all-cause and mental health and substance use-specific GP visits during the observation period. Diagnostic codes used to determine mental health and substance use-specific hospital and outpatient visits are provided in Supplemental Table 1.

2.6 Data collection

Intervention participants' demographic and administrative health care records, held by ICES, were linked using unique encoded identifiers; deterministic linkage was applied using OHIP numbers. Health care records were examined for the 12 months pre- and post-index date. The index date for exposure was the intervention enrolment date, ranging from January 7, 2013 to May 16, 2014; the study period was January 2012 to May 2015. All

participants provided written informed consent and the study received Research Ethics Board approval from Unity Health Toronto. To identify the matched cohort used in comparative analyses, population-based administrative demographic and health records were retrieved and reviewed for the same time periods.

2.7 Statistical analyses

2.7.1 Single arm pre-post analysis

Baseline characteristics for both intervention participants and individuals included in the matched cohort were calculated using frequencies and proportions for categorical variables and means and standard deviation for continuous variables. In the pre-post evaluation, inferential analyses for each count outcome included rates and rate ratios and their respective 95% confidence intervals estimated using generalized estimating equation (GEE) models with a negative binomial distribution and a log link. For each binary outcome, predicted probabilities and prevalence ratios and their respective 95% confidence intervals were estimated using GEE models with a Poisson distribution and a log link. All models were adjusted for age, sex and neighbourhood income quintile (measured at the Census tract level). An offset variable for person-years was included to adjust for different follow-up lengths.

All analyses were conducted using SAS Enterprise Guide 7.1. A significance level of .05 was used to interpret statistical significance.

2.7.2 Post-hoc comparative analysis

Baseline demographic characteristics were calculated between intervention participants with at least one hospitalization in the 12 months pre-index date and their matched counterparts using frequencies and proportions for categorical variables and means and standard deviation for continuous variables. To show trends between groups and over time, GEE models specifying a negative binomial distribution and a log link were conducted for each count outcome to estimate rate ratios and relative rate ratios; for each binary outcome, GEE models specifying a Poisson distribution and a log link were conducted to calculate prevalence ratios and relative prevalence ratios. All post-hoc models were adjusted for age, sex and neighbourhood income quintiles and included group, time and group by time interaction variables.

3. Results

The cohort selection is detailed in Figure 1. Of the 197 eligible individuals who received the intervention during the study period, 11 health records were unlinkable (six were blank and five were invalid), resulting in a cohort of 186 participants. Of those, 51

individuals were excluded from the matched analysis because they did not have a hospitalization for mental health or substance use in the 12 months prior to the index date required for comparison with the matched sample, leaving a subset of 125 individuals. Among 3,555 administrative records of people experiencing homelessness with a mental health or substance use-specific hospitalization during the exposure period, 250 individuals (1:2 matching) were selected for inclusion in the matched cohort.

3.1 Demographic characteristics

The demographic characteristics of intervention participants (n=186) are presented in Supplemental Table 2. The majority of the sample (78.5%) was male. Participants had a mean age of 40.3 years (SD=12.0), with an even distribution across age brackets between 18 and 59 years; only 4.3% of participants were over the age of 60.

Table 1 shows the demographic characteristics of the 125 intervention participants included in the matched cohort analysis and their matched counterparts; these were similar to those of the full cohort of intervention participants in age, sex, and neighbourhood income quintile as indicated by no standardized difference between the two groups being greater than 0.10.

3.2 Pre-post analysis

Rates per person-year and rate ratios (RR) for intervention participants (n=186) in the 12 months pre- and post-intervention enrolment for the number of hospital admissions, hospital days, ED visits and outpatient visits are presented in Table 2; results for binary outcomes are presented in Supplemental Table 3. Overall, results reveal changes in health care utilization patterns, with decreased inpatient care and increased outpatient care in the 12 months following enrolment in the intervention, compared to the 12 months prior. Inpatient hospital use decreased significantly among intervention participants in the 12 months following enrolment, especially for mental health and substance use-specific visits. Adjusted admission rates decreased significantly for both all cause and mental health and substance use-specific causes, by 33% (95% CI 15%-47%; $P=.001$) and 43% (95% CI 26%-66%; $P<.001$), respectively, in the 12 months post-intervention relative to 12 months prior. The total number of hospital days also decreased significantly, by 54% (95% CI 36%-77%; $P<.001$) for all-cause days and by 63% (95% CI 48%-74%; $P<.001$) for mental health and substance use days. While the number of ED visits did not change significantly over time, the overall prevalence of ED visits did decrease significantly, by 14% for any reason (95% CI 7%-20%; $P<.001$); and by 31% for mental health and substance use reasons (95% CI 12%-39%; $P<.001$) (Supplemental Table 3). In contrast to the decrease in acute hospital service use, outpatient psychiatrist visits increased by 40% (95% CI 1.15-

1
2
3
4
5
6
7
8
9 1.70; $P=.001$) in the 12 months following the intervention compared to the 12 months prior;
10
11 GP visits did not change significantly.
12

13
14 *3.3 Matched cohort analysis*
15

16
17 Rate ratios and relative rate ratios (RRR) for the subset of intervention participants
18
19 with at least one hospitalization in the 12 months prior to the index date ($n=125$) compared
20
21 to matched controls ($n=250$) 12 months pre- and post-intervention, for the number of
22
23 hospital admissions, hospital days, ED visits and outpatient visits are presented in Table 3.
24
25 Prevalence ratios (PR) and relative prevalence ratios (RPR) are presented in Supplemental
26
27 Table 4. For both count and binary outcomes, results of the matched cohort analysis suggest
28
29 that while results generally trended in the same direction as the pre-post analysis, the
30
31 changes in health service use experienced by intervention participants in the year following
32
33 receipt of the intervention as compared to the year prior were not significantly different
34
35
36
37
38 than the changes experienced by their matched counterparts (Table 3).
39
40

41 **4. Discussion**
42

43
44 Care continuity³⁵ is essential to the delivery of high quality services for people
45
46 living with chronic health conditions, including mental illness and addictions, and is
47
48 associated with improved health and service use outcomes³⁶. Our findings of decreased
49
50
51
52
53
54
55
56
57
58
59
60

acute care service use and increased outpatient service use in the 12 months following the intervention as compared to the 12 months prior are consistent with previous outcomes of critical time intervention (CTI) studies, finding significant changes in outcome measures^{24,29,30}. In addition, our findings further support and are supported by previous qualitative research suggesting improved experiences of continuity of care among adults experiencing homelessness receiving CTI or CTI adaptations^{22,37}. Notably, our findings highlight continued high rates of emergency department (ED) visits post-intervention among participants, suggesting that this population may have more acute and/or enduring needs than the brief CTI adaptation can address in our local context. Access to housing and high quality intensive case management in Toronto, for example, are extremely limited and hindered by long wait lists; it is possible that participants continued to visit the ED for immediate shelter and support post-intervention, as intervention resources were limited and not linked to housing. Persisting high ED use among participants reaffirms the need for inclusion of housing in mental health policy priorities.¹⁷

Although prior controlled studies of CTI have demonstrated improvements in acute care utilization in some settings²⁶, the addition of a matched cohort analysis in the current study indicates that changes in service use patterns of recipients of a brief CTI adaptation within our setting of universal health insurance and a relatively service-rich environment

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

are not significantly different than those of their matched counterparts. Similar to this analysis, a prior post-hoc analysis of intervention participants using self-reported data and a comparison group of adults experiencing homelessness and mental illness who were receiving usual care found successful reduction in mental health symptoms and alcohol and drug use problems among intervention participants, but increased ED visits and days spent in hospital over six months²⁴. The current post-hoc analysis, in using administrative data and a cohort matched on prior hospitalizations, was intended to build off the limitations of the aforementioned prior analysis²⁴ by focusing on service use outcomes using administrative data and exploring between-group differences with added methodological rigor. Additional measures including clinical characteristics such as diagnosis and acuity, and demographic and service use factors such as geographical location and resource availability, may help to better explain the identified non-significant differences between groups in future studies.

Our findings further highlight the need for rigorous methods in evaluating new interventions in diverse contexts. The brief CTI adaptation evaluated in the current study was set in a large urban centre under a universal health insurance system and included a dedicated, low-barrier, multidisciplinary weekly physician clinic in addition to brief case management support with limited training and supervision of frontline staff, in keeping

with a pragmatic field intervention. Our findings suggest that participants' access to preventative outpatient services was successfully facilitated by streamlined access to GPs and psychiatrists. Such an approach may be particularly relevant to similarly large urban centre where timely access to physician and case management resources is hindered by rapid general and homeless population growth and corresponding demand for services. While fidelity standards for CTI have been developed, detailing the key model components requiring adherence, and the contextual structures and staff competencies needed to ensure model integrity³⁸⁻⁴⁰, a recent systematic review indicates that fidelity and adaptations remain highly variable²⁶ and should be a required component of rigorous evaluations. The fidelity of the intervention to the CTI model was not formally assessed in this study. Future efforts should ensure that local adaptations of evidence-based interventions balance local needs and resources with fidelity to key intervention ingredients, so that evidence generated can be reliably attributed to the model of interest.

4.1 Strengths and limitations

Our findings contribute to the growing evidence on the impact of CTI and its adaptations on health service use. Results are strengthened by the study's methodological approach, which included the use of population-based administrative data and a rigorous selection process to ensure the appropriateness of comparators. However, the identification

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

of a matched cohort was challenged by the lack of a baseline acuity measure, requiring us to match on recent hospitalizations as a proxy for acuity, although post-matching demographic comparisons suggested the groups were minimally different. Additionally, intervention participants were selected using some clinical criteria that could not be applied to matched individuals identified from administrative data. In the absence of a built-in control group, this an acknowledged design limitation that could have resulted in selection bias.

We were also limited by our sample size and underpowered to detect hospitalization differences. It is possible that the lower relative rates of psychiatric hospitalizations observed in the intervention group would be significantly different with a larger sample. Still, without a randomized design, it is possible the improvements in health care use observed in the pre-post analysis are due to regression to the mean, rather than the effect of the intervention.

Given the high cost of hospitalizations and from a quality of care perspective, future research should focus on rigorous evaluations using experimental methods with large samples and robust sets of explanatory variables, in addition to approaches to strengthening fidelity of local adaptations to evidence-supported treatments. Related research should

investigate the cost-effectiveness of these interventions to maximize health outcomes given the limited resources available.

5. Conclusions

Study findings suggest that participants of a brief Critical Time Intervention (CTI) for adults experiencing homelessness and unmet mental health needs had decreased acute care use and increased outpatient service use post-intervention. Post-hoc analyses, however, found that changes in service use patterns were not significantly different from those of a matched cohort of adults experiencing homelessness. While CTI and its adaptations hold promise in improving continuity of care and health outcomes for this population, more rigorous study designs with adequate sample sizes are needed to further examine the effectiveness of CTI and local adaptations in diverse health care contexts.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Data Access

Data included in this publication are not publicly available and were collected by ICES. As a prescribed entity under Ontario’s privacy legislation, ICES is authorized to collect and use health care data for the purposes of health system analysis, evaluation and decision support. Secure access to these data is governed by policies and procedures that are approved by the Information and Privacy Commissioner of Ontario. For further questions regarding data accessibility, please contact the corresponding author.

Acknowledgements

This study was supported by ICES, which is funded by an annual grant from the Ontario Ministry of Health (MOH). Parts of this material are based on data and information compiled and provided by: MOH and the Canadian Institute for Health Information. The analyses, conclusions, opinions and statements expressed herein are solely those of the authors and do not reflect those of the funding or data sources; no endorsement is intended or should be inferred.

Conflict of Interest Disclosure: none.

Funding: This study was supported by ICES, which is funded by an annual grant from the Ontario Ministry of Health.

References

1. Abbé Pierre Foundation, European Federation of National Organisations Working with the Homeless [FEANTSA]. Fifth Overview of Housing Exclusion in Europe 2020. FEANTSA: Brussels; 2020.
2. Henry M, Watt R, Mahathey A et al. The 2019 Annual Homeless Assessment Report (AHAR) to Congress: Part 1. Point-in-time estimates of homelessness. The U.S. Department of Housing and Urban Development; 2020.
3. Gaetz S, DeJ E, Richter T et al. The State of Homelessness in Canada 2016. Toronto: Canadian Observatory on Homelessness Press; 2016.
4. Fazel S, Geddes JR, Kushel M. The health of homeless people in high-income countries: Descriptive epidemiology, health consequences, and clinical and policy recommendations. *Lancet*. 2014;384(9953):1529-1540.
[https://doi.org/10.1016/S0140-6736\(14\)61132-6](https://doi.org/10.1016/S0140-6736(14)61132-6)
5. Depp CA, Vella L, Orff HJ et al. A quantitative review of cognitive functioning in homeless adults. *J Nerv Ment Dis*. 2015;203(2):126-131.
<https://doi.org/10.1097/NMD.0000000000000248>

6. Stergiopoulos V, Cusi A, Bekele T et al. Neurocognitive impairment in a large sample of homeless adults with mental illness. *Acta Psychiatr Scand*. 2015;131(4):256-268. <https://doi.org/10.1111/acps.12391>

7. Hwang SW. Homelessness and health. *CMAJ*. 2001;164(2):229-233.

8. Aldridge RW, Story A, Hwang SW et al. Morbidity and mortality in homeless individuals, prisoners, sex workers, and individuals with substance use disorders in high-income countries: A systematic review and meta-analysis. *Lancet*. 2018;391(10117):241-250. [https://doi.org/10.1016/S0140-6736\(17\)31869-X](https://doi.org/10.1016/S0140-6736(17)31869-X)

9. Corrigan P, Pickett S, Kraus D et al. Community-based participatory research examining the health care needs of African Americans who are homeless with mental illness. *J Health Care Poor Underserved*. 2015;26(1):119-133. <https://doi.org/10.1353/hpu.2015.0018>

10. Gelberg L, Browner CH, Lejano E et al. *Women & Health*. 2004;40(2):87-100. https://doi.org/10.1300/J013v40n02_06

11. Khandor E, Mason K. *The Street Health Report 2007*. Toronto, ON: Street Health; 2007.

12. Skosireva A, O’Campo P, Zerger S et al. Different faces of discrimination: Perceived discrimination among homeless adults with mental illness in healthcare

- settings. BMC Health Serv Res. 2014;14:376. <https://doi.org/10.1186/1472-6963-14-376>
13. Buck DS, Brown CA, Mortensen K et al. Comparing homeless and domiciled patients' utilization of the Harris County, Texas public hospital system. J Health Care Poor Underserved. 2012;23(4):1660-1670. <https://doi.org/10.1353/hpu.2012.0171>
14. Fazel S, Geddes JR, Kushel M. The health of homeless people in high-income countries: Descriptive epidemiology, health consequences, and clinical and policy recommendations. Lancet. 2014;384(9953):1529-1540. [https://doi.org/10.1016/S0140-6736\(14\)61132-6](https://doi.org/10.1016/S0140-6736(14)61132-6)
15. Hwang SW, Weaver J, Aubry T et al. Hospital costs and length of stay among homeless patients admitted to medical, surgical, and psychiatric services. Med Care. 2011;49(4):350-354. <https://doi.org/10.1097/MLR.0b013e318206c50d>
16. Saab D, Bisenbaum R, Dhalla I et al. Hospital readmissions in a community-based sample of homeless adults: A matched-cohort study. J Gen Intern Med. 2016;31(9):1011-8. <https://doi.org/10.1007/s11606-016-3680-8>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

17. Pevalin DJ, Reeves A, Baker E et al. The impact of persistent poor housing conditions on mental health: A longitudinal population-based study. *Prev Med.* 2017;105:304-310. <https://doi.org/10.1016/j.ypmed.2017.09.020>

18. Klinkenberg WD, Calsyn RJ. Predictors of psychiatric hospitalization: A multivariate analysis. *Adm Policy Ment Health.* 1998;25(4):403-410. <https://doi.org/10.1023/A:1022292407868>

19. Burra TA, Hwang SW, Rourke SB et al. Homeless and housed inpatients with schizophrenia: Disparities in service access upon discharge from hospital. *Int J Ment Health Addict.* 2012;10(5):778-789. <https://doi.org/10.1007/s11469-012-9392-3>

20. Laliberte V, Stergiopoulos V, Jacob B et al. Homelessness at discharge and its impact on psychiatric readmission and physician follow-up: A population-based cohort study. *Epidemiol Psychiatr Sci.* 2020;29:e21. <https://doi.org/10.1017/S2045796019000052>

21. Hwang SW, Burns T. Health interventions for people who are homeless. *Lancet.* 2014;384(9953):1541-1547. [https://doi.org/10.1016/S0140-6736\(14\)61133-8](https://doi.org/10.1016/S0140-6736(14)61133-8)

- 1
2
3
4
5
6
7
8
9 22. Lamanna D, Stergiopoulos V, Durbin J et al. Promoting continuity of care for
10 homeless adults with unmet health needs: The role of brief interventions. *Health*
11 *Soc Care Community*. 2018;26(1):56-64. <https://doi.org/10.1111/hsc.12461>
12
13
14
15
16 23. Stergiopoulos V, Gozdzik A, Misir V et al. Effectiveness of Housing First with
17 intensive case management in an ethnically diverse sample of homeless adults with
18 mental illness: A randomized controlled trial. *PLoS One*. 2015;10(7):e0130281.
19
20
21
22 <https://doi.org/10.1371/journal.pone.0130281>
23
24
25 24. Stergiopoulos V, Gozdzik A, Nisenbaum R et al. Bridging hospital and community
26 care for homeless adults with mental health needs: Outcomes of a brief
27 interdisciplinary intervention. *Can J Psychiatry*. 2018;63(11):774-784.
28
29
30
31 <https://doi.org/10.1177/0706743718772539>
32
33
34 25. Tomita A, Herman DB. The role of a critical time intervention on the experience of
35 continuity of care among persons with severe mental illness following hospital
36 discharge. *J Nerv Ment Dis*. 2015;203(1):65-70.
37
38
39
40 <https://doi.org/10.1097/NMD.0000000000000224>
41
42
43 26. Ponka D, Agbata E, Kendall C et al. The effectiveness of case management
44 interventions for the homeless, vulnerably housed and persons with lived
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

experience: A systematic review. PLoS One. 2020;15(4):e0230896.
<https://doi.org/10.1371/journal.pone.0230896>

27. Herman DB, Conover S, Gorroochurn P et al. Randomized trial of critical time intervention to prevent homelessness after hospital discharge. Psychiatr Serv. 2011;62(7):713-719. https://doi.org/10.1176/ps.62.7.pss6207_0713

28. Herman DB, Susser ES, Jandorf L et al. Homelessness among individuals with psychotic disorders hospitalized for the first time: Findings from the Suffolk County Mental Health Project. Am J Psychiatry. 1998;155(1):109-113.
<https://doi.org/10.1176/ajp.155.1.109>

29. Shaffer SL, Hutchison SL, Ayers AM et al. Brief critical time intervention to reduce psychiatric rehospitalization. Psychiatr Serv. 2015;66(11):1155-1161.
<https://doi.org/10.1176/appi.ps.201400362>

30. Tomita A, Herman DB. The impact of critical time intervention in reducing psychiatric rehospitalization after hospital discharge. Psychiatr Serv. 2012;63(9):935-937. <https://doi.org/10.1176/appi.ps.201100468>

31. KasproW WJ, Rosenheck RA. Outcomes of critical time intervention case management of homeless veterans after psychiatric hospitalization. Psychiatr Serv. 2007;58(7):929-935. <https://doi.org/10.1176/ps.2007.58.7.929>.

32. Jones K, Colson PW, Holter MC. Cost-effectiveness of critical time intervention to reduce homelessness among persons with mental illness. *Psychiatr Serv*. 2003;54(6):884-90. <https://doi.org/10.1176/appi.ps.54.6.884>
33. Stergiopoulos V, Gozdzik A, Nisenbaum R et al. Integrating hospital and community care for homeless people with unmet mental health needs: Program rationale, study protocol and sample description of a brief multidisciplinary case management intervention. *Int J Ment Health Addict*. 2017;15:362-378. <https://doi.org/10.1007/s11469-017-9731-5>
34. Austin PC. Using the standardized difference to compare the prevalence of a binary variable between two groups in observational research. *Commun Stat-Simul Comput*. 2009;38(6):1228-1234. <https://doi.org/10.1080/03610910902859574>
35. Haggerty JL, Reid RJ, Freeman GK et al. Continuity of care: A multidisciplinary review. *Br Med J*. 2003;327(7425):1219-1221. <https://doi.org/10.1136/bmj.327.7425.1219>
36. Puntis S, Rugkasa J, Forrest A et al. Associations between continuity of care and patient outcomes in mental health care: A systematic review. *Psychiatr Serv*. 2015;66(4):354-363. <https://doi.org/10.1176/appi.ps.201400178>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

37. Dixon L, Goldberg R, Iannone V et al. Use of a critical time intervention to promote continuity of care after psychiatric inpatient hospitalization. *Psychiatr Serv.* 2009;60(4):451-458. <https://doi.org/10.1176/ps.2009.60.4.451>

38. da Silva TFC, Lovisi GM, Conover S. Developing an instrument for assessing fidelity to the intervention in the Critical Time Intervention – Task Shifting (CTI-TS) – preliminary report. *Arch Psychiatry Psychother.* 2014;1:55-62. <https://doi.org/10.12740/APP/23279>

39. De Vet R, Lako DAM, Jeijersbergen MD et al. Critical time intervention for people leaving shelters in the Netherlands: Assessing fidelity and exploring facilitators and barriers. *Adm Policy Ment Health.* 2017;44:67-80. <https://doi.org/10.1007/s10488-015-0699-9>

40. Shaw J, Conover S, Herman D et al. Critical time intervention for severely mentally ill prisoners (CrISP): A randomized controlled trial. *Health Services and Delivery Research.* 2017;5(8). <https://doi.org/10.3310/hsdr05080>

Tables

Table 1: Demographic characteristics of CATCH-H participants *with at least one hospitalization in the 12 months pre-intervention* (n=125) and matched controls (n= 250)

Demographic variables		CATCH-H participants with hospitalization (n=125)		Matched controls (n=250)		Standardized difference
		n	%	n	%	
Age (years)	18 to 29	29	23.2	58	23.2	0
	30 to 39	27	21.6	52	20.8	0.02
	40 to 49	36	28.8	73	29.2	0.01
	50 to 59	25	20	49	19.6	0.01
	60 and over	8	6.4	18	7.2	0.03
	Mean (SD)	41.1 (12.4)		41	12.3	0
Sex	Female	27	21.6	52	20.8	0
	Male	98	78.4	198	79.2	
Neighbourhood income quintile	1* (low)	39	31.2	90	36	0.09
	2 (medium low)	30	24	55	22	0.05
	3 (medium)	26	20.8	49	19.6	0.03
	4 (medium high)	13	10.4	23	9.2	0.04
	5 (high)	17	13.6	33	13.2	0.01

Legend: SD = standard deviation

*1: includes individuals in the 1st income quintile and those where this information was missing

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2: Rates per person-year and rate ratios (RR) for CATCH-H participants (n=186) 12 months pre-intervention vs. 12 months post-intervention for number of hospital admissions, hospital days, ED visits and outpatient visits estimated from adjusted generalized estimating equations with negative binomial distribution

Outcome variable		CATCH-H participants (n=186)						
		Rate per person-year				Rate ratio (RR)		
		12 months pre-intervention		12 months post-intervention		RR	95% CI	P-value
		Rate	95% CI	Rate	95% CI			
Hospital admissions	All cause	2.00	1.54-2.60	1.35	0.96-1.88	0.67	0.53-0.85	.001
	Mental health and substance use	1.22	0.92-1.63	0.69	0.48-1.01	0.57	0.44-0.74	<.001
Days in hospital	All cause	24.66	18.93-32.12	11.37	8.09-15.99	0.46	0.33-0.64	<.001
	Mental health and substance use	17.67	12.34-25.32	6.47	4.22-9.93	0.37	0.26-0.52	<.001
ED visits	All cause	8.98	6.76-11.94	9.57	6.90-13.29	1.07	0.90-1.25	.446
	Mental health and substance use	3.77	2.79-5.08	3.57	2.49-5.11	0.95	0.75-1.19	.645
Outpatient visits	Psychiatrist	3.16	2.36-4.25	4.42	3.32-5.88	1.40	1.15-1.70	.001
	GP	7.49	6.30-8.91	7.70	6.63-8.94	1.03	0.88-1.20	.725
		3.38	2.70-4.23	3.80	3.16-4.58	1.13	0.93-1.37	.224
	Total outpatient visits	11.05	9.35-13.07	12.64	10.81-14.77	1.14	1.00-1.31	.049

(Psychiatrist +
GP)

Legend: CI = confidence interval; GP = general practitioner

Table 3: Rate ratios (RR) and relative rate ratios (RRR) for *CATCH-H* participants with at least one hospitalization in the 12 months pre-intervention (n=125) compared to matched controls (n=250) 12 months pre-intervention vs. 12 months post-intervention for number of hospital admissions, hospital days, ED visits and outpatient visits estimated using adjusted generalized estimating equations with negative binomial distribution


Outcome variable		12 months pre-intervention			12 months post-intervention			Relative rate ratio (RRR) 12 months post-intervention vs. 12 months pre-intervention		
		RR	95% CI	P-value	RR	95% CI	P-value	RR	95% CI	P-value
Hospital admissions	All cause	1.20	0.99-1.45	.057	1.17	0.82-1.66	.38	0.97	0.71-1.33	.870
	Mental health and substance use	1.06	0.87-1.29	.565	0.95	0.64-1.39	.78	0.89	0.63-1.26	.522
Days in hospital	All cause	1.04	0.75-1.43	.808	0.77	0.51-1.17	.22	0.74	0.46-1.20	.225
	Mental health and substance use	0.93	0.67-1.29	.667	0.60	0.38-0.95	.03	0.64	0.38-1.10	.104
ED visits	All cause	1.35	1.01-1.79	.040	1.69	1.19-2.42	.004	1.26	0.95-1.67	.113

Outpatient visits	G P	Mental health and substance use	0.95 -	1.33	1.86	.096	1.64	1.11-2.43	.01	1.23	0.89-1.71	.212
		Psychiatrist	0.96 -	1.29	1.74	.091	1.34	1.02-1.77	.04	1.04	0.77-1.40	.806
		All cause	0.96 -	1.22	1.56	.102	1.36	1.06-1.75	.02	1.11	0.86-1.44	.409
		Mental health and substance use	0.85 -	1.16	1.58	.356	1.32	0.98-1.77	.07	1.14	0.84-1.55	.408
		Total outpatient visits (Psychiatrist + GP)	1.02 -	1.24	1.52	.035	1.33	1.10-1.62	.003	1.07	0.87-1.33	.522

Legend: CI = confidence interval; GP = general practitioner

Figures

Figure 1. Exposed and matched cohort selection



Supplementary Materials

Supplemental Table 1: Diagnostic codes used in cohort selection

Mental health outcomes	Diagnostic Codes
Hospitalizations	ICD-10-CA: all F codes (main diagnosis) OMHRS: all discharges
Emergency department visits	ICD-10-CA: F04 to F99 (main diagnosis) ICD-10-CA: X60-X84, Y10-Y19, Y28 (any diagnosis)
Psychiatrist visits	OHIP visit/consult to a psychiatrist [SPEC = 19]
General practitioner outpatient visits	OHIP visit/consult to a general practitioner [SPEC = 00] AND a mental health diagnostic code:295, 296, 297, 298, 300, 301, 302, 306, 309, 311, 303, 304, 897, 898, 899, 900, 901, 902, 904, 905, 906,909

Supplemental Table 2: Demographics of CATCH-H cohort (n=186)

Demographic variables		N	%
Age (years)	18 to 29	43	23.1
	30 to 39	46	24.7
	40 to 49	51	27.4
	50 to 59	38	20.4
	60 and over	8	4.3
	Mean (SD)	40.3 (12.0)	
Sex	Female	40	21.5
	Male	146	78.5
Neighbourhood income quintile	Missing	6	3.2
	1 (low)	56	30.1
	2 (medium low)	44	23.7
	3 (medium)	41	22.0
	4 (medium high)	16	8.6
	5 (high)	23	12.4

Legend: SD = standard deviation

Supplemental Table 3: Predicted probabilities and prevalence ratios (PR) for CATCH-H participants (n=186) 12 months pre-intervention vs. 12 months post-intervention for any hospital admission, any ED visit, and any outpatient visit estimated from adjusted generalized estimating equations with Poisson distribution

Outcome variable		CATCH-H participants (n=186)						
		Predicted probability (PP)				Prevalence ratio (PR)		
		12 months pre-intervention		12 months post-intervention		PR	95% CI	P-value
		PP	95% CI	PP	95% CI			
Any hospital admission	All cause	0.77	0.68-0.86	0.48	0.40-0.57	0.62	0.52-0.74	<.001
	Mental health	0.58	0.46-0.73	0.31	0.24-0.41	0.54	0.44-0.67	<.001
Any ED visit	All cause	0.96	0.91-1.01	0.83	0.77-0.89	0.86	0.80-0.93	<.001
	Mental health	0.79	0.71-0.87	0.54	0.47-0.63	0.69	0.61-0.78	<.001
Any outpatient visit	Psychiatrist	0.74	0.65-0.85	0.69	0.60-0.79	0.93	0.84-1.03	.174
	GP	0.86	0.80-0.93	0.89	0.83-0.95	1.03	0.96-1.10	.392
	Mental health	0.73	0.65-0.80	0.76	0.68-0.84	1.04	0.94-1.16	.449
	Total outpatient visits (Psychiatrist + GP)	0.95	0.92-0.99	0.98	0.95-1.01	1.03	0.98-1.07	.231

Legend: CI = confidence interval; GP = general practitioner

Supplemental Table 4: Prevalence ratios (PR) and relative prevalence ratios (RPR) for *CATCH-H* participants with at least one hospitalization in the 12 months pre-intervention (n=125) compared to matched controls (n=250) 12 months pre-intervention vs. 12 months post-intervention for any hospital admission, ED visit and outpatient visit estimated using adjusted generalized estimating equations with Poisson distribution

Outcome variable		12 months pre-intervention			12 months post-intervention			Relative prevalence ratio (RPR)		
		PR	95% CI	P-value	PR	95% CI	P-value	RP R	95% CI	P-value
Any hospital admission	All cause	1.00	0.99-1.01	.982	0.89	0.72-1.10	.30	0.89	0.72-1.10	.297
	Mental health	0.86	0.80-0.92	<.001	0.78	0.60-1.00	.047	0.91	0.71-1.17	.450
Any ED visit	All cause	1.03	0.99-1.06	.142	1.07	0.96-1.20	.20	1.05	0.94-1.17	.428
	Mental health	0.97	0.89-1.05	.456	0.91	0.75-1.11	.36	0.94	0.78-1.14	.546
Any outpatient visit	Psychiatrist	1.13	1.01-1.26	.040	1.09	0.96-1.25	.18	0.97	0.83-1.14	.728
	GP	1.08	0.98-1.19	.120	1.16	1.06-1.27	.002	1.07	0.96-1.20	.239
			0.99-1.32	.066	1.26	1.10-1.45	.001	1.10	0.93-1.31	.259
	Total outpatient visits (Psychiatrist + GP)	1.03	0.97-1.09	.368	1.03	0.98-1.09	.24	1.01	0.93-1.08	.890

Legend: CI = confidence interval; GP = general practitioner

Évaluer l’effet d’une adaptation de l’intervention en temps critique sur l’utilisation des soins de santé chez des adultes itinérants ayant des besoins de santé mentale dans un grand centre urbain

Objectif : La présente étude a évalué l’effet d’une adaptation de l’intervention en temps critique (ITC) sur les résultats de l’utilisation des soins de santé chez des adultes en situation d’itinérance et ayant des besoins de santé mentale dans un grand centre urbain.

Méthode : Des données administratives provinciales dans la population de l’Ontario, Canada, ont servi à une méthode avant et après pour une cohorte de 197 personnes qui ont reçu l’intervention entre janvier 2013 et mai 2014, et ont été appariées à une cohorte d’adultes en situation d’itinérance qui n’ont pas reçu l’intervention durant la même période. Les changements des résultats de l’utilisation des soins de santé dans l’année avant et après l’intervention ont été évalués à l’aide d’équations d’estimation généralisées et des analyses a posteriori ont évalué les différences entre les groupes.

Résultats : Les analyses avant et après ont révélé des changements statistiquement significatifs des modèles d’utilisation des soins de santé chez les bénéficiaires de l’intervention, notamment une utilisation réduite des services pour les patients hospitalisés et une utilisation accrue des services pour les patients ambulatoires dans l’année suivant l’intervention comparé à l’année précédente. Toutefois, l’analyse de la cohorte appariée a constaté des différences non significatives dans les changements d’utilisation des services de santé entre un sous-groupe de bénéficiaires de l’intervention et leurs homologues appariés.

Conclusions : Un modèle adapté d’ITC était associé à des changements de l’utilisation des soins de santé chez les personnes en situation d’itinérance et ayant des besoins de santé mentale. Cependant, les changements n’étaient pas différents de ceux observés dans une cohorte appariée. Il faut des méthodes rigoureuses et des échantillons adéquats dans les études qui examinent l’efficacité de l’ITC et les adaptations locales dans divers contextes de soins de santé.